



Nanotechnology application in overcoming the reproductive disorders in livestock

Muhammad Wasif Gulzar*¹, Muhammad Kasib Khan¹, Rimsha Gulzar¹, Muhammad Suleman¹, Jawad Hussain¹, Ahmad Hassan¹, Muhammad Abdullah², Ahmad Raza², Zulqarnain Haider¹

¹ University of Agriculture, Faisalabad, Pakistan

² Govt. Willayat Hussain Islamia College, Multan, Pakistan

Article info

Received: 01 May 2024

Received in revised form: 10 June 2024

Accepted: 06 July 2024

Published online: 20 July 2024

Keywords

Nano-delivery system
Reproductive disorders
Biostimulation
Nutrition
Hormones
Antibiotics

* Corresponding author:

Muhammad Wasif Gulzar

Email: 2022ag5900@uaf.edu.pk

Reviewed by:

Consent to publish the name of reviewers could not be obtained

Abstract

The profitability as well as productivity of livestock farming operations are significantly impacted by the reproduction rate of farm animals. Applying several techniques, such as dietary, hormonal, and biological methods together with controlling reproductive diseases, is the foundation of optimal reproductive management. These tactics ought to adhere to ethical and practical standards in addition to providing adequate reproductive results. For instance, a number of biological parameters and an animal's physiological state are primarily related to the effectiveness of biological and hormonally based reproductive tactics. Additional aspects, such as digestion and absorption, may also play a role in the effectiveness of nutritional strategies. Concerns about the overuse of antibiotics or the emergence of antibiotic-resistant bacteria further complicate the management of illnesses connected to reproduction. The use of nanotechnology in fields such as cattle farming systems could offer novel and creative ways to address problems with reproductive control. Nanotechnology can give several pharmaceuticals (including hormones and antibiotics), molecules from biology, and nutrients with new physicochemical qualities. These include enhanced bioavailability, increased cellular absorption, regulated sustained release, and decreased toxicity as compared to conventional versions. In this review, it will be demonstrated how nanotechnology has advanced the most popular reproductive management systems while taking into account the ongoing difficulties associated with each strategy.

This is an open access article under the CC Attribution-ND license (<http://creativecommons.org/licenses/by-nd/4.0/>)

1. Introduction

Reproductive efficiency in cattle farming systems has a direct impact on farm output, profitability, and sustainability. The effectiveness of producing milk and/or meat is determined by the reproduction rate of farm animals. This can be done directly or indirectly through management choices like replacement and harvesting rates. Applying precise techniques is the cornerstone of optimal reproductive management, which also requires taking costs, animal welfare, environmental effects, and human health into account. Following the selection of a strategy that aligns with each farm's objectives, the majority of reproductive management techniques are prepared for use in economic livestock farms (Olynk and Wolf 2008; Smith et al. 2018). These tactics could involve nutritional management, reproductive aided techniques (mostly artificial insemination and estrous synchronisation), a number of bio stimulating instruments (like the male effect), and the prevention or treatment of reproductive illnesses (Caraviello et al. 2006; Delgadillo and Martin 2015; Hassanein et al. 2021). Despite being widely and primarily used in various livestock production systems, the effectiveness of these reproductive management techniques is

called into question by a number of pragmatic and moral considerations. For instance, hormone-based reproductive medicines are the recommended approach to managing reproduction; yet, the pharmacokinetics and pharmacodynamics of these treatments are critical to their efficacy and can be influenced by biological factors (Hashem and Gonzalez-Bulnes 2020; Hassanein et al. 2021). The male effect is a method of sexual biostimulation that offers the chance to do away with the heavy reliance on hormones for reproductive management and the creation of hormone-free animal products. Agriculture is another industry that has seen a transformation because of artificial insemination (AI). AI has significantly improved cattle raising in particular in a variety of ways. This technology has the potential to raise animal welfare standards, boost sustainable farming practices, and increase output (Gulzar and Hussain 2023). Nevertheless, the male to female ratio, age, and experience of the male all pose challenges to the method's effectiveness (Chasles et al. 2016; Tejada et al. 2017; Ungerfeld 2007). Similarly, inadequate delivery of necessary nutrients and low nutrient bioavailability may have a negative effect on nutritional management strategies aimed at enhancing farm animal reproductive performance (Hashem

and El-Zarkouny 2014; Izquierdo et al. 2015). Last but not least, worries about the widespread use of antibiotics and the emergence of bacterial resistant to antibiotics pose a barrier to the treatment of illnesses connected to reproduction (Algharib et al. 2020; Cerbu et al. 2021; Yang et al. 2009). Taking these factors into account, the development of new technologies, such as nanotechnology, opens up a world of possibilities for livestock and agricultural output.

Nano-drug delivery systems represent the most significant and promising use of nanotechnologies in the animal production industry. By utilising nanotechnology, a variety of pharmaceuticals, biological molecules, and nutrients can gain new physicochemical characteristics, including enhanced bioavailability, greater mobility and uptake by cells, regulated sustained distribution of the medicine at the target site, decreased toxicity in comparison to other compounds, enhanced enzymatic actions, and elevated mucoadhesive properties (Osama et al. 2020). With the use of nanotechnology, this review aims to demonstrate potential advancements in the most popular reproductive management techniques.

2. Hormone based treatments

Even with current efforts to control reproduction in animals used for agriculture using biological methods, the practices—including exogenous hormone therapy, cannot be completely eliminated from the farming sector. Treatments based on hormones are a useful tool for increasing fertility and the profitability of farming operations. Numerous survey investigations supported the importance of hormonal-based procedures for reproductive management; for example, 87% (103 of 153) of managers from large milking herds (average herd size of 613 cows) in the USA confirmed the use of hormonal synchronisation or timed artificial insemination in their reproductive management (Caraviello et al. 2006). Similarly, about 80% of practitioners from 714 inorganic dairy farms in England confirmed the crucial role of hormones for efficient reproductive management (Higgins et al. 2013). Hormones are frequently utilised because they are necessary for the implementation of many assisted reproductive techniques such as timed artificial insemination (Hashem et al. 2015), superovulation, and estrous synchronisation/induction (Hashem et al. 2015) for increasing the efficiency of reproduction and for treating disorders related to reproduction in both males and females (Hashem et al. 2015; Hashem and Aboul-ezz 2018). Progesterone, oestrogen, testosterone, melatonin, prostaglandins, and gonadotropins are the primary hormones used to regulate fertility in farm animals. The pharmacokinetics and pharmacodynamics of these hormones are primarily responsible for their efficacy. Certain hormones, like prostaglandin F₂ α (PGF₂ α) and gonadotropin-releasing hormone (GnRH), have short half-lives and low molecular weights, which limits their ability to be delivered to target areas over time. Certain hormones, like glycoprotein gonadotropins (follicle-stimulating hormone – FSH; luteinizing hormone – LH; human chorionic gonadotropin – hCG; equine chorionic gonadotropin – eCG), can trigger the immune system and the production of particular antibodies, which can make some farm animal species resistant to repeated gonadotropic treatments

(Castro et al. 2009). Additionally, reduced fertility and other biological consequences are linked to recurrent administration of these hormones. Anti-eCG antibodies have been demonstrated to impede eCG bioactivities through two different methods. The first method involves blocking eCG's interaction with its receptors; the second involves anti-eCG antibodies changing eCG's structure, which can block eCG bioactivities. It is important to note that after repeated treatments, fertility is impacted since these modifications of eCG biological activity by its antibodies mostly affect the FSH bioactivity of eCG (Herve et al. 2004; Kara et al. 2019).

In this instance, ewes receiving up to three rounds of eCG/FSH based super ovulatory treatments displayed decreased rates of fertilisation and less viable embryos overall at the subsequent recoveries in comparison to the first flushing (Forcada et al. 2011). In a different study, the kidding rate (41.3%) was lower in goats with high eCG levels of antibodies following repeated eCG treatments than in other females (66.7%). The preovulatory LH surge and delayed estrous rate were linked to these goats' reduced fertility (Roy et al. 1999). Similar results were seen in rabbits, where repeated administration of recombinant human FSH elevated the levels of FSH antibodies in the females at the time of the third and fourth superovulation treatments. Concerns about animal health and welfare, the environment, and managing reproduction and fertility in farm animals present additional difficulties to the usage of hormones. For instance, since eCG is derived from bleeding pregnant mares, a shortage is anticipated owing to concerns about animal welfare. Future hormone production may be halted by the persistent social pressure placed on corporations producing the hormone (Manteca Vilanova et al. 2019). Lastly, because hormone residues and carrier materials are released into the environment, conventional hormonal delivery methods may also throw off the equilibrium of the surrounding ecosystem. The employing of progesterone-impregnated intravaginal gadgets which were created to regulate the estrous cycle in various agricultural animals, is the most obvious example (Rathbone and Burke 2013). The progesterone-loaded silicon polymers that make up the majority of these devices must have high concentrations of progesterone in order to release enough hormones to the genital area mucosa. This raises the possibility of hormone emissions into the environment, direct hormone transmission to workers or breeders, and indirect hormone transmission to shoppers through animal products (de Graaff and Grimard 2018).

3. Reproduction related diseases

Reproductive illnesses are typically linked to the phase of animal production, and more especially, reproduction (Casares-Crespo et al. 2018; Fernandez-Serrano et al. 2017; Hashem and Sallam 2020). Reduced conception rate and an increased risk of reproductive culling are two associated symptoms of postpartum diseases, particularly endometritis caused by various bacterial species (primarily *Escherichia coli*, *Staphylococcus aureus*, *Bacillus cereus*, *Pseudomonas aeruginosa*, *Prevotella melaninogenica*, *Arcanobacterium pyogenes*) (Helbling et al. 2014; Oliveira et al. 2013; Rather et al. 2013). When compared

to their healthy herd mates, the risk of pregnancy and the number of days from calving until pregnancy were 40 %; HR = 0.60; 69 %; HR = 0.31; 76 %; HR = 0.24 in cows with metritis, clinical endometritis, as well as subclinical endometritis, respectively (Fogolari et al. 2016; Remiao et al. 2016). Protozoan illnesses linked to pregnancy-associated disorders, like toxoplasmosis (*Toxoplasma gondii*) and neosporosis (*Neospora caninum*), cause large financial losses in farm ruminants. Globally, neosporosis infections is a prevalent zoonosis (Li et al. 2018; Mahdasht et al. 2020) that results in abortion in cattle, while toxoplasmosis infection primarily produces reproductive dysfunction in small ruminants (Helbling et al. 2018). Bovine mastitis, primarily caused by *Staphylococcus aureus*, is a major source of economic losses in dairy farms. The disease results in steep drops in milk yield (about 380 tons of milk are lost globally annually), tainted milk, reproductive disorders, and additional costs for replacing tainted milk and replacing infected animals (Cordova-Izquierdo 2016; Santos-Jimenez et al. 2020). Furthermore, *Staphylococcus aureus* contamination of raw milk poses a threat to public health along the entire food chain (Cannas et al. 2019; Zhao et al. 2021). In general, these illnesses have detrimental effects on population health, animal welfare, and the ultimate profit from the manufacturing process (Scaramuzzi and Martin 2008; Wu 2010). The majority of these bacterium and/or zoonotic diseases include clinical symptoms that are either directly caused by tissue damage and microbial products (endotoxin) or indirectly caused by mediators of oxidant stress (nitric oxide) and/or inflammation (cytokines and eicosanoids) (Penagaricano et al. 2013).

Antibiotic-based therapy is currently the most widely advised course of treatment for a variety of microbial/protozoan diseases, including illnesses connected to reproduction (Castro et al. 2019; Sturmeijer et al. 2009). The metabolism of the medication regulates the efficacy of antibiotic-based therapy. The rate at which antibiotics are absorbed and distributed determines whether they will reach the intended infected sites or not (Hippen et al. 2008; Hosny et al. 2020; Shin et al. 2012; Zereu 2016). This rate can be influenced by a variety of biological factors, including the antibiotics' resistance to gastrointestinal enzyme degradation when administered orally, blood hydrolytic enzymes when administered parenterally, drug solubility, and consequent cellular absorption and bioavailability. Furthermore, some infections cause fibrous damage in the infected tissues, which limits the reach of the medications into infected regions when regional therapy is administered—for example, when a medication is directly infused into the female reproductive organ in cases of endometritis or administered through the teats in cases of mastitis (Boerman and Lock 2014; El-Sherbiny et al. 2016; Gawad and Fellner 2019; Hackmann and Firkins 2015; Hammon et al. 2005). It is important to consider the risks associated with the emergence of new wild pathogenic microbial species, the spread of infectious and cross-transmitted microbial species, the accidental spillage of antibiotics into the environment, and the transfer of antibiotic residues into livestock products (meat and milk) (Jahanbin et al. 2015). In addition of being detrimental to the general public's health, antimicrobial resistance increases the chance of

treatment failure, recurrent infection, and related economic consequences (Khalil et al. 2019). In fact, because of these variables, using antibiotics to treat disorders associated with reproduction (such as mastitis, toxoplasmosis, and neosporosis) is a contentious approach. Therefore, new, secure, and efficient antibiotic therapy based strategies are required, especially when treating farm animals that are used for food production (Albuquerque et al. 2020; Shahin et al. 2020).

4. Nanotechnology approaches

Numerous studies have demonstrated the potential to overcome the therapeutic limitations of antibiotic-based therapies (Gurunathan et al. 2018; Vallejo-Timaran et al. 2020) by employing a variety of engineered nanomaterials (such as liposomes, nanoparticles of polymers, solid lipid nanoparticles, nanogels, and inorganic nanoparticles) that are synthesised with specific physicochemical properties (Sanchez-Sanchez et al. 2018; Zhou et al. 2018). The use of nano-formula for antibiotic-based therapies may also reduce the dosage of the antibiotic, enable effective delivery of the medication to the infected sites, shorten the duration of therapy and minimise side effects and antibiotic degradation (Olsen et al. 2006; Piotr et al. 2013). Nanomaterials have the potential to enhance the transport of antibiotics to diseased sites and provide protection against their rapid degradation (Wang et al. 2017). Additionally, nanomaterials can be made to exhibit cytotoxic and destructive capabilities against microbes.

Furthermore, certain nanoparticles cause harm to the bacterial cell membrane, enzymes, and structural and functional proteins of the cell primarily by inducing oxidative pathways in the cell that alter gene expression and promote both innate and adaptive immunity. It is possible to tailor nanoparticles so that they prevent bacterial adhesion, colonisation, and biofilm formation (Algharib et al. 2020). Additionally, pharmaceuticals can be incorporated into nanostructures without changing the compound's structure, thereby enhancing its pharmacological efficacy (Gholipourmalekabadi et al. 2017). In particular, the antibiotic enrofloxacin is used to treat a variety of bacterial illnesses in pigs, including *Salmonella*, *Pasteurella*, *Mycoplasma*, and *Escherichia coli*. The suggested intramuscular dose for swine is 2.5 to 5 mg of enrofloxacin/kg Bw/day for three to five days. It has been demonstrated that drinking water with a suspension of enrofloxacin-loaded poly (lactic-co-glycolic acid) nanostructures could be administered orally, and that this would result in a 23% decrease in the minimum inhibitory concentration against *E. coli* when compared to enrofloxacin alone (El-Zawawy et al. 2015; Paudel et al. 2019). In a different investigation, atovaquone nanosuspensions coated with sodium dodecyl sulphate significantly improved the transit through gastrointestinal and blood–brain barriers, hence increasing the therapeutic efficacy against experimentally acquired and reactivated toxoplasmosis (Shubar et al. 2011). Similarly, for *Staphylococcus aureus* lactation infection, tilmicosin (a semi-synthetic macrolide antibiotic) - loaded hydrogenated castor oil at a decreased dosage has shown superior therapeutic efficacy than free tilmicosin due to its greater bioavailability and sustained-release performance (Wang et al. 2012). In order

to address intracellular persistence of *Staphylococcus aureus* and multi-drug resistance, which are linked to the subclinical and recurrent infection of bovine mastitis, nanoparticle medicines have also been employed as a tactic recently. Yang demonstrated how the use of amoxicillin nanoparticles in the treatment of bovine mastitis may extend the duration of the post-antibiotic effects and, consequently, the intervals between doses (Yang et al. 2009).

Innovative healthy and secure antibiotic alternatives are becoming possible due to the recent convergence of the benefits of nano-drug delivery methods and alternative medicine, which relies on the use of natural compounds with antibacterial activity. Levamisole and morantel are two examples of imidazothiazole medications that have been widely used to treat *Haemonchus contortus* infections; however, the development of resistance has called into question their continued efficacy (Qamar and Alkheraije 2023). Probiotic species, microbial extracts from plants, and botanical secondary metabolites (polyphenols and essential oils) have all been identified in numerous studies as possible antibacterial agents (Garzon et al. 2021). In a different investigation, the use of poly (lactic-co glycolic) acid (PLGA)-epigallocatechin gallate-doxycycline nanoparticles as an aided endometritis therapy proved beneficial (Garzon et al. 2021). In this case, mastitis has been treated with chitosan-TPP nanoparticles (Rivera Aguayo et al. 2020). Moreover, certain metal nanoparticles, such as copper oxide (CuO), zinc oxide (ZnO), silver oxide (Ag₂O), gold (Au), and titanium dioxide (TiO₂) have shown strong antibacterial activity against a variety of microorganisms. These alternatives may offer a chance for patients to completely replace out antibiotic-based medications for safer ones (Ferreira-Silva and Burnett 2017). Because nanometals (silver) can be synthesized biologically while meeting environmental and therapeutic standards, the use of nanomaterials as an antibiotic substitute has been promoted. The anthelmintic closantel belongs to the class of medications known as salicylanilide. Target blood-sucking parasites such the *Haemonchus* species have plasma proteins that salicylanilide binds to with a strong affinity and high specificity. Moreover, Closantel may interfere with the parasite's defense systems to preserve pH imbalance (Qamar and Alkheraije 2023). The natural reducing agents such as flavonoids, polyphenols, and other flavonoids and phenolic acids molecules of life of Camellia, green tea, when black tea leaf extracts are used, along with microorganisms like *E. Coli*, a type of *Acinetobacter*, *Pseudomonas aeruginosa*, and *Klebsiella pneumoniae* for the nanoparticle synthesis (Das et al. 2020). In this case, the manufacture of silver nanoparticles measuring 10 nm was accomplished through the use of apigenin, a polyphenolic molecule. By preventing the viability of cells and creation of biofilm in a dose-and time-dependent manner, these nanoparticles demonstrated antibacterial efficacy against pathogenic bacteria *Prevotella melaninogenica* and *Arcanobacterium pyogenes* found in an endometrial infected uterine discharges.

Similarly, Yuan verified that two triple drug-resistant strains of *Pseudomonas aeruginosa* and *Staphylococcus aureus*

obtained from goat milk samples afflicted with mastitis could be inhibited by biologically synthesised silver nanoparticles (Yuan et al. 2017). Radzikowski confirmed the ability of readily available silver nanoparticles, metal nanoparticles, and the combination of them to reduce the viability about mastitis-borne pathogenic organisms without exhibiting harmful impacts on mammary gland tissues, in reference to the toxicity of such nanotechnologies to animal tissues (Radzikowski et al. 2020). Additionally, Paudel verified that the drug's integration into the PLGA matrix reduced the amount of reactive oxygen compounds that the antibiotic induced, indicating that enrofloxacin-entrapped nanoparticles are less harmful to mammalian cells than the free drug (Paudel et al. 2019).

5. Conclusions

This review highlighted the current drawbacks and restrictions of the most widely used reproductive management techniques in farm animals. As demonstrated, factors limiting the effectiveness of reproductive management strategies include ease of field-scale application, animal behaviour and physiological state, drug availability and uptake, and environmental factors like antibiotic/hormone residue release. Innovative and unconventional solutions are provided by nanotechnology. Nanotechnology has the potential to limit the emergence of antibiotic-resistant microbial species by increasing the effectiveness of antibiotics and/or developing natural antibiotic substitutes. However, in light of the information now accessible, this study offers bold remedies for the problems with reproductive management; further research is needed to determine the efficacy of these tactics.

Declarations

Funding: This review article received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflict of interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements: None

References

- Albuquerque J, Casal S, Pascoa RNM de J, Van Dorpe I, Fonseca AJM, Cabrita ARJ, Neves AR, and Reis S. (2020). Applying nanotechnology to increase the rumen protection of amino acids in dairy cows. *Scientific Reports* 10(1): 6830. <https://doi.org/10.1038/s41598-020-63793-z>
- Algharib SA, Dawood A, Xie S. (2020). Nanoparticles for treatment of bovine *Staphylococcus aureus* mastitis. *Drug Delivery* 27(1): 292–308. <https://doi.org/10.1080/10717544.2020.1724209>
- Boerman JP, Lock AL. (2014). Effect of unsaturated fatty acids and triglycerides from soybeans on milk fat synthesis and biohydrogenation intermediates in dairy cattle. *Journal of Dairy Science* 97(11): 7031–7042. <https://doi.org/10.3168/jds.2014-7966>
- Cannas A, Tedeschi LO, Atzori AS, Lunesu MF. (2019). How can nutrition models increase the production efficiency of sheep and goat operations? *Animal Frontiers* 9(2): 33–44. <https://doi.org/10.1093/af/vfz005>

- Caraviello DZ, Weigel KA, Fricke PM, Wiltbank MC, Florent MJ, Cook NB, Nordlund KV, Zwald NR, Rawson CL. (2006). Survey of management practices on reproductive performance of dairy cattle on large US commercial farms. *Journal of Dairy Science* 89(12): 4723–4735. [https://doi.org/10.3168/jds.s0022-0302\(06\)72522-x](https://doi.org/10.3168/jds.s0022-0302(06)72522-x)
- Casares-Crespo L, Fernandez-Serrano P, Viudes-de-Castro MP. (2018). Protection of GnRH analogue by chitosan-dextran sulfate nanoparticles for intravaginal application in rabbit artificial insemination. *Theriogenology* 116: 49–52. <https://doi.org/10.1016/j.theriogenology.2018.05.008>
- Castro MV, Cortell C, Moce E, Marco-Jimenez F, Joly T, Vicente J, Forcada F, Ait Amer-Meziane M, Abecia JA, Maurel MC, Cebrian-Perez JA, Muino-Blanco T, Asenjo B, Vazquez MI, Casao A. (2009). Effect of recombinant gonadotropins on embryo quality in superovulated rabbit does an immune response after repeated treatments. *Theriogenology* 72: 655–662. <https://doi.org/10.1016/j.theriogenology.2009.04.022>
- Castro T, Martinez D, Isabel B, Cabezas A, Jimeno V. (2019). Vegetable oils rich in polyunsaturated fatty acids supplementation of dairy cows' diets: Effects on productive and reproductive performance. *Animals* 9(5): 205. <https://doi.org/10.3390/ani9050205>
- Cerbu C, Kah M, White JC, Astete CE, Sabliov CM. (2021). Fate of biodegradable engineered nanoparticles used in veterinary medicine as delivery systems from a one health perspective. *Molecules* 26: 523. <https://doi.org/10.3390/molecules26030523>
- Chasles M, Chesneau D, Moussu C, Delgadillo JA, Chemineau P, Keller M. (2016). Sexually active bucks are efficient to stimulate female ovulatory activity during the anestrus season also under temperate latitudes. *Animal Reproduction Science* 168: 86–91. <https://doi.org/10.1016/j.anireprosci.2016.02.030>
- Cordova-Izquierdo A. (2016). Best practices in animal reproduction: Impact of nutrition on reproductive performance livestock. *Journal of Advances in Dairy Research* 4(1): 152. <https://doi.org/10.4172/2329-888X.1000152>
- Das CA, Kumar VG, Dhas TS, Karthick V, Govindaraju K, Joselin JM, Baalamurugan J. (2020). Antibacterial activity of silver nanoparticles (biosynthesis): A short review on recent advances. *Biocatalysis and Agricultural Biotechnology* 27: 101593. <https://doi.org/10.1016/j.bcab.2020.101593>
- de Graaff W, Grimard B. (2018). Progesterone-releasing devices for cattle estrus induction and synchronization: Device optimization to anticipate shorter treatment durations and new device developments. *Theriogenology* 112: 34–43. <https://doi.org/10.1016/j.theriogenology.2017.09.025>
- Delgadillo JA, Martin GB. (2015). Alternative methods for control of reproduction in small ruminants: A focus on the needs of grazing industries. *Animal Frontiers* 5(1): 57–65. <https://doi.org/10.2527/af.2015-0009>
- El-Sherbiny M, Cieslak A, Pers-Kamczyc E, Szczechowiak J, Kowalczyk D, Szumacher-Strabel M. (2016). A nanoemulsified form of oil blends positively affect the fatty acid proportion in ruminal batch cultures. *Journal of Dairy Science* 99: 399–407. <https://doi.org/10.3168/jds.2015-9328>
- El-Zawawy LA, El-Said D, Mossallam SF, Ramadan HS, Younis SS. (2015). Triclosan and triclosan-loaded liposomal nanoparticles in the treatment of acute experimental toxoplasmosis. *Experimental Parasitology* 149: 54–64. <https://doi.org/10.1016/j.exppara.2014.12.007>
- Fernandez-Serrano P, Casares-Crespo L, Viudes-de-Castro M. (2017). Chitosan–dextran sulphate nanoparticles for GnRH release in rabbit insemination extenders. *Reproduction in Domestic Animals* 52(S4): 72–74. <https://doi.org/10.1111/rda.13062>
- Ferreira-Silva JC, Burnett. (2017). Luteinizing hormone (LH) levels and ovarian activity in postpartum Santa Inês ewes are subject to a male effect. *Archives of Animal Breeding* 60(2): 95–100. <https://doi.org/10.5194/aab-60-95-2017>
- Fogolari O, Felipe AC, Leimann FV, Goncalves OH, Sayer C, Araujo PHD. (2016). Method validation for progesterone determination in poly (Methyl methacrylate) nanoparticles synthesised via miniemulsion polymerization. *International Journal of Polymer Science* 2017(1): 9603140. <https://doi.org/10.1155/2017/9603140>
- Forcada F, Ait Amer-Meziane M, Abecia JA, Maurel MC, Cebrian-Perez JA, Muino-Blanco T, Asenjo B, Vazquez MI, Casao A. (2011). Repeated superovulation using a simplified FSH/eCG treatment for in vivo embryo production in sheep. *Theriogenology* 75(4): 769–776. <https://doi.org/10.1016/j.theriogenology.2010.10.019>
- Garzon S, Lagana AS, Barra F, Casarin J, Cromi A, Raffaelli R, Uccella S, Franchi M, Ghezzi F, Ferrero S. (2021). Novel drug delivery methods for improving efficacy of endometriosis treatments. *Expert Opinion on Drug Delivery* 18(3): 355–367. <https://doi.org/10.1080/17425247.2021.1829589>
- Gawad R, Fellner V. (2019). Evaluation of glycerol encapsulated with alginate and alginate-chitosan polymers in gut environment and its resistance to rumen microbial degradation. *Asian Australasian Journal of Animal Science* 32(1): 72–81. <https://doi.org/10.5713/ajas.18.0110>
- Gholipourmalekabadi M, Mobaraki M, Ghaffari M, Zarebkohan A, Omrani VF, Urbanska AM, Seifalian A. (2017). Targeted drug delivery based on gold nanoparticle derivatives. *Current Pharmaceutical Design* 23(20): 2918–2929. <https://doi.org/10.2174/1381612823666170419105413>
- Gulzar MW, Hussain J. (2023). The implementation of artificial intelligence in veterinary diseases and practices: A comprehensive review. *Hosts and Viruses* 10: 43–50. <https://doi.org/10.17582/journal.hv/2023/10.43.50>
- Gurunathan S, Choi Y-J, Kim J-H. (2018). Antibacterial efficacy of silver nanoparticles on endometritis caused by *Prevotella melanogenica* and *Arcanobacterium pyogenes* in dairy cattle. *International Journal of Molecular Sciences* 19(4): 1210. <https://doi.org/10.3390/ijms19041210>
- Hackmann TJ, Firkins JL. (2015). Maximising efficiency of rumen microbial protein production. *Frontiers in Microbiology* 6: 137653. <https://doi.org/10.3389/fmicb.2015.00465>
- Hammon DS, Holyoak GR, Dhiman TR. (2005). Association between blood plasma urea nitrogen levels and reproductive fluid urea nitrogen and ammonia concentrations in early lactation dairy cows. *Animal Reproduction Science* 86(3): 195–204. <https://doi.org/10.1016/j.anireprosci.2004.08.003>
- Hashem NM, Aboul-ezz ZR. (2018). Effects of a single administration of different gonadotropins on day 7 post-insemination on pregnancy outcomes of rabbit does. *Theriogenology* 105: 1–6. <https://doi.org/10.1016/j.theriogenology.2017.09.006>
- Hashem NM, El-Azrak KM, Nour El-Din ANM, Taha TA, Salem MH. (2015). Effect of GnRH treatment on ovarian activity and reproductive performance of low-prolific Rahmani ewes. *Theriogenology* 83(2): 192–198. <https://doi.org/10.1016/j.theriogenology.2014.09.016>

- Hashem NM, El-Zarkouny S, Taha T, Abo-Elezz Z. (2015). Oestrous response and characterization of the ovulatory wave following oestrous synchronisation using PGF 2α alone or combined with GnRH in ewes. *Small Ruminant Research* 129: 84–87. <https://doi.org/10.1016/j.smallrumres.2015.06.003>
- Hashem NM, El-Zarkouny SZ. (2014). Effect of short-term supplementation with rumen-protected fat during the late luteal phase on reproduction and metabolism of ewes. *Journal of Animal Physiology and Animal Nutrition* 98(1): 65–71. <https://doi.org/10.1111/jpn.12032>
- Hashem NM, Gonzalez-Bulnes A. (2020). State-of-the-art and prospective of nanotechnologies for smart reproductive management of farm animals. *Animals* 10(5): 840. <https://doi.org/10.3390/ani10050840>
- Hashem NM, Sallam S. (2020). Reproductive performance of goats treated with free gonadorelin or nano conjugated gonadorelin at estrus. *Domestic Animal Endocrinology* 71: 106390. <https://doi.org/10.1016/j.domaniend.2019.106390>
- Hassanein E, Hashem N, El-Azrak K, Gonzalez-Bulnes A, Hassan G, Salem M. (2021). Efficiency of GnRH-loaded chitosan nanoparticles for inducing LH secretion and fertile ovulations in protocols for artificial insemination in rabbit does. *Animals* 11: 440. <https://doi.org/10.3390/ani11020440>
- Helbling IM, Busatto CA, Fioramonti SA, Pesoa JI, Santiago L, Estenez DA, Luna JA. (2018). Preparation of TPP-crosslinked chitosan microparticles by spray drying for the controlled delivery of progesterone intended for estrus synchronisation in cattle. *Pharmaceutical Research* 35(3): 66. <https://doi.org/10.1007/s11095-018-2363-z>
- Helbling IM, Ibarra JCD, Luna JA. (2014). The optimization of an intravaginal ring releasing progesterone using a mathematical model. *Pharmaceutical Research* 31(3): 795–808. <https://doi.org/10.1007/s11095-013-1201-6>
- Herve V, Roy F, Bertin J, Guillou F, Maurel MC. (2004). Anti- Equine chorionic gonadotropin (eCG) antibodies generated in goats treated with eCG for the induction of ovulation modulate the luteinizing hormone and follicle-stimulating hormone bioactivities of eCG differently. *Endocrinology* 145: 294–303. <https://doi.org/2020071619250817100>
- Higgins HM, Ferguson E, Smith RE, Green MJ. (2013). Using hormones to manage dairy cow fertility: The clinical and ethical beliefs of veterinary practitioners. *PLOS ONE* 8(4): e62993. <https://doi.org/10.1371/journal.pone.0062993>
- Hippen AR, DeFrain JM, Linke PL. (2008). Glycerol and other energy sources for metabolism and production of transition dairy cows. *Proceedings of the Florida Ruminant Nutrition Symposium*. January 29-30, Best Western Gateway Grand, Gainesville, Florida, USA.
- Hosny NS, Hashem NM, Morsy AS, Abo-elezz ZR. (2020). Effects of organic selenium on the physiological response, blood metabolites, redox status, semen quality, and fertility of rabbit bucks kept under natural heat stress conditions. *Frontiers in Veterinary Science* 7: 529584. <https://doi.org/10.3389/fvets.2020.00290>
- Izquierdo AC, Gutierrez JFP, Hernandez WM, Mancera AEV, Crispín RH. (2015). Obtencion, evaluación y manipulación del semen de verraco en una unidad de producción mexicana. *Revista Veterinaria* 26(1): 69-74. <https://doi.org/10.30972/vet.261253>
- Jahanbin R, Yazdanshenas P, Amin Afshar M, Mohammadi Sangcheshmeh A, Varnaseri H, Chamani M, Nazaran MH, Bakhtiarizadeh MR. (2015). Effect of zinc nano-complex on bull semen quality after freeze-thawing process. *Animal Production* 17(2): 371-380. <https://doi.org/10.22059/jap.2015.54040>
- Kara E, Dupuy L, Bouillon C, Casteret S, Maurel MC. (2019). Modulation of gonadotropins activity by antibodies. *Frontiers in Endocrinology* 10: 440002. <https://doi.org/10.3389/fendo.2019.00015>
- Khalil WA, El-Harairy MA, Zeidan AEB, Hassan MAE. (2019). Impact of selenium nano-particles in semen extender on bull sperm quality after cryopreservation. *Theriogenology* 126: 121–127. <https://doi.org/10.1016/j.theriogenology.2018.12.017>
- Li Y, Zhao X, Wang L, Liu Y, Wu W, Zhong C, Zhang Q, Yang J. (2018). Preparation, characterization and in vitro evaluation of melatonin-loaded porous starch for enhanced bioavailability. *Carbohydrate Polymers* 202: 125–133. <https://doi.org/10.1016/j.carbpol.2018.08.127>
- Manteca Vilanova X, De Briyne N, Beaver B, Turner PV. (2019). Horse welfare during equine chorionic gonadotropin (eCG) production. *Animals* 9(12): 1053. <https://doi.org/10.3390/ani9121053>
- Oliveira JE, Medeiros ES, Cardozo L, Voll F, Madureira EH, Mattoso LH, Assis OB. (2013). Development of poly (lactic acid) nanostructured membranes for the controlled delivery of progesterone to livestock animals. *Materials Science & Engineering C, Materials for Biological Applications* 33(2): 844–849. <https://doi.org/10.1016/j.msec.2012.10.032>
- Oliveira JE, Medeiros ESC, L., V. F., M., E.H., M., L.H.C., A., & O.B.G. (2013). Development of poly (lactic acid) nanostructured membranes for the controlled delivery of progesterone to livestock animals. *Mater. Sci. Eng. C*, 33,844-849.
- Olsen JE, Christensen H, Aarestrup FM. (2006). Diversity and evolution of bla Z from *Staphylococcus aureus* and coagulase-negative staphylococci. *Journal of Antimicrobial Chemotherapy* 57(3): 450–460. <https://doi.org/10.1093/jac/dki492>
- Olynk N, Wolf C. (2008). Economic analysis of reproductive management strategies on US commercial dairy farms. *Journal of Dairy Science* 91(10): 4082-4091.
- Osama E, El-Sheikh SM, Khairy MH, Galal AA. (2020). Nanoparticles and their potential applications in veterinary medicine. *Journal of Advanced Veterinary Research* 10: 268–273.
- Paudel S, Cerbu C, Astete CE, Louie SM, Sabliov C, Rodrigues DF. (2019). Enrofloxacin-impregnated PLGA nanocarriers for efficient therapeutics and diminished generation of reactive oxygen species. *ACS Applied Nano Materials* 2(8): 5035–5043. <https://doi.org/10.1021/acsanm.9b00970>
- Penagaricano F, Souza AH, Carvalho PD, Driver AM, Gamba R, Kropp J, Hackbart KS, Luchini D, Shaver RD, Wiltbank MC. (2013). Effect of maternal methionine supplementation on the transcriptome of bovine preimplantation embryos. *PLoS ONE* 8: 72302. <https://doi.org/10.1371/journal.pone.0072302>
- Piotr S, Marta S, Aneta F, Barbara K, Magdalena Z. (2013). Antibiotic resistance in *Staphylococcus aureus* strains isolated from cows with mastitis in eastern Poland and analysis of susceptibility of resistant strains to alternative non-antibiotic agents: Lysostaphin, nisin and polymyxin B. *Journal of Veterinary Medical Science* 76(3): 355-362. <https://doi.org/10.1292/jvms.13-0177>
- Qamar W, Alkheraije KA (2023). Anthelmintic resistance in *Haemonchus contortus* of Sheep and Goats from Asia-a review of in vitro and in vivo studies. *Pakistan Veterinary Journal* 43(3): 376–387.

- Radzikowski D, Kalinska A, Ostaszewska U, Gołębiewski M. (2020). Alternative solutions to antibiotics in mastitis treatment for dairy cows—a review. *Animal Science Papers and Reports* 38(2): 117–133.
- Rathbone MJ, Burke CR. (2013). Controlled release intravaginal veterinary drug delivery. In: MJ Rathbone, A McDowell, editors, *Long acting animal health drug products: Fundamentals and applications*, Springer, USA. pp. 247–270.
https://doi.org/10.1007/978-1-4614-4439-8_11
- Rather MA, Sharma R, Gupta S, Ferosekhan S, Ramya V, Jadhao SB. (2013). Chitosan-nano conjugated hormone nanoparticles for sustained surge of gonadotropins and enhanced reproductive output in female fish. *PLoS One* 8: 57094.
<https://doi.org/10.1371/journal.pone.0057094>
- Remiao MH, Lucas CG, Domingues WB, Silveira T, Barther NN, Komninou ER, Basso AC, Jornada DS, Beck RCR, Pohlmann AR, Junior ASV, Seixas FK, Campos VF, Guterres SS, Collares T. (2016). Melatonin delivery by nanocapsules during in vitro bovine oocyte maturation decreased the reactive oxygen species of oocytes and embryos. *Reproductive Toxicology* 63: 70–81.
<https://doi.org/10.1016/j.reprotox.2016.05.016>
- Rivera Aguayo P, Bruna Larenas T, Alarcon Godoy C, Cayupe Rivas B, Gonzalez-Casanova J, Rojas-Gomez D, Caro Fuentes N. (2020). Antimicrobial and antibiofilm capacity of chitosan nanoparticles against wild type strain of *Pseudomonas* sp. isolated from milk of cows diagnosed with bovine mastitis. *Antibiotics* 9(9): 551.
<https://doi.org/10.3390/antibiotics9090551>
- Roy F, Maurel M-C, Combes B, Vaiman D, Crihiu EP, Lantier I, Pobel T, Deletang F, Combarous Y, Guillou F. (1999). The negative effect of repeated equine chorionic gonadotropin treatment on subsequent fertility in alpine goats is due to a humoral immune response involving the major histocompatibility complex1. *Biology of Reproduction* 60(4): 805–813.
<https://doi.org/10.1095/biolreprod60.4.805>
- Sanchez-Sanchez R, Vazquez P, Ferre I, Ortega-Mora LM. (2018). Treatment of toxoplasmosis and neosporosis in farm ruminants: State of knowledge and future trends. *Current Topics in Medicinal Chemistry* 18(15) 1304–1323.
<https://doi.org/10.2174/1568026618666181002113617>
- Santos-Jimenez Z, Guillen-Gargallo S, Encinas T, Berlinguer F, Veliz-Deras FG, Martinez-Ros P, Gonzalez-Bulnes A. (2020). Use of propylene-glycol as a cosolvent for GnRH in synchronization of estrus and ovulation in sheep. *Animals* 10(5): 897.
<https://doi.org/10.3390/ani10050897>
- Scaramuzzi R, Martin G. (2008). The importance of interactions among nutrition, seasonality and socio-sexual factors in the development of hormone-free methods for controlling fertility. *Reproduction in Domestic Animals* 43(S2): 129–136.
<https://doi.org/10.1111/j.1439-0531.2008.01152.x>
- Shahin MA, Khalil WA, Saadeldin IM, Swelum AAA, El-Harairy MA. (2020). Comparison between the effects of adding vitamins, trace elements, and nanoparticles to SHOTOR extender on the cryopreservation of dromedary camel epididymal spermatozoa. *Animals* 10(1): 78.
<https://doi.org/10.3390/ani10010078>
- Shin JH, Wang D, Kim SC, Adesogan AT, Staples CR. (2012). Effects of feeding crude glycerin on performance and ruminal kinetics of lactating Holstein cows fed corn silage- or cottonseed hull-based, low-fiber diets. *Journal of Dairy Science* 95(7): 4006–4016.
<https://doi.org/10.3168/jds.2011-5121>
- Shubar HM, Lachenmaier S, Heimesaat MM, Lohman U, Mauludin R, Mueller RH, Fitzner R, Borner K, Liesenfeld O. (2011). SDS-coated atovaquone nanosuspensions show improved therapeutic efficacy against experimental acquired and reactivated toxoplasmosis by improving passage of gastrointestinal and blood–brain barriers. *Journal of Drug Targeting* 19(2): 114–124.
<https://doi.org/10.3109/10611861003733995>
- Siahdasht FN, Farhadian N, Karimi M, Hafizi L. (2020). Enhanced delivery of melatonin loaded nanostructured lipid carriers during in vitro fertilisation: NLC formulation, optimization and IVF efficacy. *RSC Advances* 10: 9462–9475.
<https://doi.org/10.1039/c9ra10867j>
- Smith ME, Geisert RD, Parrish JJ. (2018). Reproduction in domestic ruminants during the past 50 years: Discovery to application. *Journal of Animal Science* 96(7): 2952–2970.
<https://doi.org/10.1093/jas/sky139>
- Stewart F, Allen W, Moor RM. (1976). Pregnant mare serum gonadotropin: Ratio of follicle-stimulating hormone and luteinizing hormone activities measured by radioreceptor assay. *Journal of Endocrinology* 71: 371–382.
<https://doi.org/10.1677/joe.0.0710371>
- Sturmey R, Reis A, Leese H, McEvoy T. (2009). Role of fatty acids in energy provision during oocyte maturation and early embryo development. *Reproduction in Domestic Animals* 44(S3): 50–58.
<https://doi.org/10.1111/j.1439-0531.2009.01402.x>
- Tejada LM, Meza-Herrera CA, Rivas-Munoz R, Rodriguez-Martínez R, Carrillo E, Mellado M, Veliz-Deras FG. (2017). Appetitive and consummatory sexual behaviors of rams treated with exogenous testosterone and exposed to anestrus dorper ewes: Efficacy of the male effect. *Archives of Sexual Behavior* 46(3): 835–842.
<https://doi.org/10.1007/s10508-016-0852-x>
- Ungerfeld R. (2007). Socio-sexual signalling and gonadal function: Opportunities for reproductive management in domestic ruminants. *Society of Reproduction and Fertility Supplement* 64: 207–221.
<https://doi.org/10.5661/rdr-vi-207>
- Vallejo-Timaran DA, Arango-Sabogal JC, Reyes-Vélez J, Maldonado-Estrada JG. (2020). Postpartum uterine diseases negatively impact the time to pregnancy in grazing dairy cows from high-altitude tropical herds. *Preventive Veterinary Medicine* 185: 105202.
<https://doi.org/10.1016/j.prevetmed.2020.105202>
- Wang L, Hu C, Shao L. (2017). The antimicrobial activity of nanoparticles: Present situation and prospects for the future. *International Journal of Nanomedicine* 2017(12): 1227–1249.
<https://doi.org/10.2147/IJN.S121956>
- Wang XF, Zhang SL, Zhu LY, Xie SY, Dong Z, Wang Y, Zhou WZ. (2012). Enhancement of antibacterial activity of tilmicosin against *Staphylococcus aureus* by solid lipid nanoparticles in vitro and in vivo. *The Veterinary Journal* 191(1): 115–120.
<https://doi.org/10.1016/j.tvjl.2010.11.019>
- Wu G. (2010). Functional amino acids in growth, reproduction, and health. *Advances in Nutrition* 1(1): 31–37.
<https://doi.org/10.3945/an.110.1008>
- Yang X, Ouyang W, Sun J, Li X. (2009). Post-antibiotic effect of Amoxicillin nanoparticles against main pathogenic bacteria of bovine mastitis in vitro. *Journal of Northwest Agriculture and Forestry University (Natural Science Edition)* 37: 1–6.
- Yuan YG, Peng QL, Gurunathan S. (2017). Effects of silver nanoparticles on multiple drug-resistant strains of *Staphylococcus aureus* and *Pseudomonas aeruginosa* from mastitis-infected goats: An alternative approach for antimicrobial therapy. *International Journal of Molecular Sciences* 18(3): 569.

<https://doi.org/10.3390/ijms18030569>

Zereu G. (2016). Factors affecting feed intake and its regulation mechanisms in ruminants a review. *International Journal of Livestock Research* 6: 19–40.

<https://doi.org/10.5455/ijlr.20160328085909>

Zhao C, Bai Y, Fu S, Wu L, Xu C, Xia C. (2021). Follicular fluid proteomic profiling of dairy cows with anestrus caused by negative energy balance. *Italian Journal of Animal Science* 20(1): 650–663.

<https://doi.org/10.1080/1828051x.2021.1899855>

Zhou K, Li C, Chen D, Pan Y, Tao Y, Qu W, Liu Z, Wang X, Xie S. (2018). A review on nanosystems as an effective approach against infections of *Staphylococcus aureus*. *International Journal of Nanomedicine* 13: 7333–7347. <https://doi.org/10.2147/ijn.s169935>

Citation

Gulzar MW, Khan MK, Gulzar R, Suleman M, Hussain J, Hassan A, Abdullah M, Raza A, Haider Z. (2024). Nanotechnology application in overcoming the reproductive disorders in livestock. *Letters in Animal Biology* 04(2): 11 – 18.